



Correction to: Effectiveness of fissure sealants following different silver fluoride application protocols in MIH-affected molars: randomized clinical study

Nagihan Cayiroglu¹ · Elif Ballikaya¹ · Gizem Erbas Unverdi¹ · Zafer Cavit Cehreli²

© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2026

Correction to: Clin Oral Invest

<https://doi.org/10.1007/s00784-026-06856-1>

In the published paper, the authors identified an error in the naming of the study groups in the Materials and Methods section.

The group definitions were incorrectly reported. As presented in the tables and the main text, Groups 1, 3, and 5 correspond to the resin sealant groups. The corrected group names are provided below, and this correction has no impact on the analyses, results, or conclusions of the study.

The corrected group names are as follows:

Group 1: Resin-based fissure sealant (RBS; 3M™ Clinpro™ Sealant, 3 M ESPE, St. Paul, MN, USA), applied one week after SF application.

Group 2: Glass ionomer fissure sealant (GIS; GC Fuji TRIAGE®, GC Europe N.V., Leuven, Belgium), applied one week after SF application.

Group 3: RBS applied immediately after SF application following blot-drying.

Group 4: GIS applied immediately after SF application following blot-drying.

Group 5: RBS applied immediately after SF application following rinsing.

Group 6: GIS applied immediately after SF application following rinsing.

The original article has been corrected.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s00784-026-06856-1>.

✉ Elif Ballikaya
eyildirim@hacettepe.edu.tr

Nagihan Cayiroglu
nagihan.nc27@gmail.com

Gizem Erbas Unverdi
erbasgizem@gmail.com

Zafer Cavit Cehreli
zcehreli@gmail.com

¹ Department of Pediatric Dentistry, Faculty of Dentistry, Hacettepe University, Ankara 06100, Turkey

² Department of Pediatric Dentistry, Faculty of Dentistry, Istanbul Health and Technology University, Istanbul, Turkey